

THE UNIVERSITY OF ARIZONA.

College of Agriculture and Life Sciences

AZ1435

11/08

Ten Steps to a Successful Vegetable Garden

Gardening with vegetables can be fun and provide delicious and highly nutritious fresh food. Watching and working with plants can add a new dimension of enjoyment to life and bring an awareness of the wonderful world of nature in the backyard. The marvels of nature will have special personal meaning when nurturing a small seed into a colorful productive plant with your own hands. These accomplishments can be obtained regardless of the size of garden. A few plants or a large plot will give rewarding experiences for both young and old. The path to a successful vegetable garden is not difficult or long. Ten carefully taken steps will produce many enjoyable moments and an abundant harvest of fresh vegetables during much of the year.

Select a good location

Choose an area with plenty of morning sunlight and some afternoon shade. Most vegetables, especially fruiting types, do best with six to eight hours of full sun exposure. Leafy and root vegetables will tolerate partial shade. Don't plant gardens under or near trees or large shrubs—their roots will rob fertility and water from vegetables. Don't plant vegetables in the narrow shaded space between houses and walls.

A loose, fertile, level, well-drained soil is best. If possible, avoid heavy clays and very sandy soils. If caliche is present it must be dug out and removed. Avoid areas that are crusted with alkali salts or infested with Bermudagrass, nutgrass or Johnson grass.

A synthetic soil, self prepared or purchased, can be used in raised beds or containters (pots, tubs, boxes) if good soil is not available. Where the space is limited, container gardening can be practiced. A convenient water supply for irrigating is necessary.

Plan your garden layout

Planning ahead will help avoid problems and make your garden a complement to your landscape. First, sketch a plan of the intended planting area for vegetables. Write down the size of the area or location of containers. This is the beginning of a gardening notebook or journal.

• Decide on the vegetable species wanted. Select those that your household likes, that are adapted to your climate and practical for the location. If space is a problem, plant those that utilize space efficiently like bush varieties or bush beans, beets, broccoli, cabbage, carrots, leaf lettuce, onions, radishes, Swiss chard, tomatoes, and turnips.

 Mark on the plan where the vegetables will be planted, making sure to leave room for growing space between plants. Also, list the planting date for each vegetable. Arrange plantings according to harvest periods and growth characteristics. Plant vegetables adjacent to each other which will be harvested about the same time. Avoid having taller plants shade younger and smaller vegetables. Use vertical space by trellising climbing crops.

Grow recommended varieties

Step 3

Gardening success can be greatly influenced by the varieties you use. Select from recommended lists and from those know to do well locally. It is a good idea to try one or two new varieties each year. Plant them next to old favorites for comparison. Keep a notebook or journal from year to year to note what varieties perform best. For mini-gardens, try bush or dwarf varieties and the more colorful ones. Seed catalogues will be a big help in finding these. Look for All-American Selection Award winners. (www.all-americaselections.org/)

Step

N

Obtain good seed, plants, equipment and supplies

Before planting, find a reputable source for seed and other garden supplies. Seed catalogs can be a big help, but be sure the varieties are locally adapted. Buy new seed since some seeds over a year old will not germinate (sprout) well. Some seeds can be saved and are best placed in jars or in plastic bags and stored in a freezer.

Step

Vegetable transplants can be purchased at garden stores, nurseries and greenhouses. Insist on recommended varieties. Select plants that are healthy, stocky, mediumsized, with vigorous roots and that are pest free. Avoid plants that are wilted, yellow, spindly, too large or have spots on the leaves, brown lesions on the stems or knots/ galls on the roots. Obtain plants in containers (pots, 6 or 8 packs, bands or boxes) when possible so that the root systems are intact. Transplants should not be disturbed any more than necessary and should be "hardened-off." Transplants can be started at home if desired. Have all equipment and tools clean and in good condition before working the soil. A hoe, spade, garden rake, trowel, measuring stick and planting line are essential. A hand cultivator and seed drill reduce work in larger gardens. Hoses and sprinklers convenient for watering are also needed. Other needed supplies are fertilizers and mulching materials.

Study pest control recommendations to determine what may be needed after positively identifying the pest. It is important to have a quick source of materials for pest control if needed. A good sprayer or duster to control garden pests should be available for use. Care should be taken in handling, applying and storing all chemicals. Always follow the pesticide label instructions; it is a legal document!

Prepare and care for the soil properly

Soil provides nutrients and water for plants. If these materials are limited or if the soil is compact or hard and crusty when dry, and water-soaked and sticky when wet, plants will not grow and develop properly. To maintain and improve soil conditions, mix organic matter and fertilizers into the soil before planting, and prepare and cultivate the soil when dry or slightly moist (never when wet).

Organic matter makes the soil loose (friable) and easy to work. It improves nutrient and water-holding capacity, drainage and aeration. Well rotted manure, compost, and leaf mulch are commonly used organic materials. Composted manure is easy to use and is relatively free of weed seeds. Apply a layer of organic matter 2 to 3 inches thick on the garden area about 1 to 2 months before planting. Work it into the top 10-12 inches of soil. A thorough watering of soil at this time helps leach harmful salts from the root zone. If poultry manures are used, apply them at half rate.

A fertilizer should be added containing both nitrogen and phosphorus and be applied before planting. These nutrients will benefit most garden crops. Although soils vary in fertility, a typical fertilizer application would be 1 to 2 lbs. (1 to 2 cups) of 16-20-0 (ammonium phosphate) per 100 ft.² spread evenly over the soil. Also, 3 to 5 lb. of soil sulfur/100 ft.² may be added if water drainage is poor. All these materials should be plowed, roto-tilled or spaded into the top 10 to 12 inches of soil shortly before planting.

In preparing the seedbed, do not work the soil when it is too wet. Wait for it to dry sufficiently so it crumbles in your hands. Level the area by raking. Then make raised beds if using furrow irrigation (*See Figure A*). Top dress planted area with a three inch layer of organic mulch after seedlings emerge or after transplanting (*See Step 8*).

When growing vegetables in close quarters or where good soil is not available, an artificial soil can be used. If the soil doesn't drain well consider using raised beds filled with ½ garden soil and ½ artificial soil mix, coarse sand, perlite or vermiculite. (*see Figure B*).

During the growing season fertilizers may be needed. Applying bands of fertilizer, usually only nitrogen, is called "side-dressing." Apply ½ lb./100 feet of row of 21-0-0 or equivalent fertilizer, three inches deep and about four inches to the side of the plants. Alternatively, spread nitrogen fertilizer on the soil surface about 4 inches from the plant and water it in. However, too much fertilizer too close to the plant may injure plant roots. Examples of side-dressing timing are: tomatoes—after the first clusters of tomatoes form; sweet corn—when plants are "knee high" and again when they tassel; cucumbers, melons and squash when they begin to produce runners.

Plant your vegetables properly

Most vegetables are started from seed or transplants. Seed can be sown directly into the garden soil, while transplants are started elsewhere and later planted into the garden. Harvest can be obtained sooner with transplants; however, it is more expensive and certain plants do not transplant well. Generally, beans, beets, carrots, cucumbers, lettuce, muskmelons, onions, peas, pumpkin, radish, spinach, squash, sweet corn and watermelon are started in

6

the garden from seed. Vegetables like asparagus, broccoli, cabbage, cauliflower, eggplant, peppers, sweet potatoes and tomatoes are generally transplanted, but care needs to be taken to minimize root drying and injury.

A few simple rules need to be followed in seeding:

- Mark out straight rows to make the garden attractive and to make cultivation, insect control and harvesting easier. To mark a row, drive two stakes into the ground at each end of the garden and draw a string tightly between them. Shallow furrows, suitable for small seed, can be made by drawing a hoe handle along the line indicated by the string. For deeper furrows, use the corner of the hoe blade. Use correct spacing between rows.
- Space seeds properly in the row. The number of seeds to sow per foot or hill (more than one seed/ hole) is suggested on seed packages or in reference materials. Space the seeds uniformly. Sometimes small seeds can be handled better if they are mixed with dry, pulverized soil or sand and then spread. To aid in spacing seed, spread on one layer of toilet paper placed on the soil. The contrast of the white toilet paper will aid in seeing seed spacing.
- Plant at the proper depth. A general rule to follow is to place the seed at a depth about four times the diameter of the seed. Cover small seeds such as carrots and lettuce with no more than ¼ to ½ inch of soil. Place large seeds such as corn, beans and peas 1 to 2 inches deep. In sandy soils plant seed somewhat deeper.
- Cover seeds and firm the soil over them by gently tamping the soil by hand or the flat back of a hoe. This prevents rain or sprinkler water from washing away the seeds.
- Irrigate by sprinkling the soil surface lightly. When using furrow irrigation, hold water until moisture moves across seed row. Seeds need moisture to

germinate. Water often enough to prevent crusting and drying around the seed. After plants emerge, water less often but deeper.

• Thin plants to the desired number as soon as possible. Remove weaker plants. Scissors can aid in thinning by cutting out young plants. Do not wait too long before thinning or injury will result from crowding and disturbing the remaining plants.

When transplanting follow these directions:

- Transplant on a cloudy day or in the evening.
- Handle plants with care. About an hour before transplanting thoroughly water plants and soil in the containers (pots, bands, flats). Carefully remove plants from their containers, disturbing the roots as little as possible. Try to keep the "soil ball" around the roots. Keep roots moist at all times when they are out of the soil. If roots are "pot bound" tease them out before planting.
- Dig a hole large enough so that the transplanted plant sits slightly deeper than it grew in the container.
- Use a starter solution to get plants off to a faster start. Starter fertilizer is a soluble fertilizer high in phosphorous like 10-52-17 or 10-50-10 mixture. Mix fertilizer with water following the label directions (about 2 tablespoons per gallon of water). After plants are set in the soil, pour about 1 cup of solution around the roots of each plant. When peat or fiber pots are set in the soil add enough water to soften pot. Also, break off any excessive pot material so it is below the garden soil level to prevent water wicking. Remove any plastic or wooden bands from around roots.
- Cover the roots with soil and firm the soil around the plant.
- Protect plants for a few days from sun, wind or cold if necessary.

Irrigate with care

Irrigation is necessary for all garden crops in Arizona because of limited and uncertain rainfall. Water enough to keep the soil moist (not wet) in the root zone of the plant throughout the growing season. Excessive fluctuations of soil moisture adversely affect plant growth and quality. Regular applications of water need to be made to prevent the soil from becoming too dry (*see Figure C*).

Proper watering can be accomplished by observing the plant and soil. Do not allow the plant to become stressed, wilted or slow-growing. On the other hand, too much water, especially on heavy soils, will exclude air from the root zone, resulting in poor growth. When the soil becomes crumbly upon squeezing, it's time to irrigate. Moisture is needed around the seed for sprouting. Frequent watering will be needed to keep the soil adequately moist and prevent crusting of the surface. A three inch layer of organic mulch will help prevent evaporation. Do not place mulch on top of seedlings or transplants, but around them.

As the plant grows, the watering period should be longer, allowing deeper penetration through the root zone. Determine the moisture depth with a spade or by probing with a stick, trowel or iron rod. Most vegetables are shallow-rooted and use water from the upper 12 to 24 inches of soil.

Frequency of watering depends on many things. A large plant needs more water than a small plant. A shallow-rooted vegetable (cabbage, onion, lettuce, corn)

Step 6

Step 7

needs to be irrigated more often than a deep-rooted vegetable (asparagus, tomato, watermelon). Coarse textured soils (sandy loams) need to be irrigated more often than fine-textured (clay or silt loams). Plants need to be watered more often during hot periods than cool periods. In an average situation during warm weather, a good soaking of the soil every 5 to 7 days should give satisfactory results with established plants.

The following irrigation methods are commonly used: furrow, sprinkler, soaker hoses and drip (trickle). The furrow method delivers water alongside the plant row. Water should be kept in the furrow long enough for moisture to completely infiltrate the soil of the root zone. Garden sprinklers apply water on both plants and soil and should not be used if the water is salty. Drip or trickle emitter systems and soaker hoses apply water through a hose which lies beside the crop row. All four methods have a place in Arizona gardens. Traditionally, a raised bed with two rows is used with furrow irrigation, while a flat bed with no furrows is normally used with the other methods. If a watering method moistens the plant foliage, irrigate in the morning so plants have time to dry during the day. This will lessen disease problems. Night time watering encourages disease growth.

Plants growing in containers should be watched more closely for water needs because the roots are more crowded and temperatures of root media are more extreme. Keep soil moist but do not over-water. Make holes on the side and/or the bottom of the container for drainage and air.

Mulch & cultivate to control weeds

Weeds compete with vegetables for water, nutrients and light. Weeds often harbor insects and diseases. Two important ways to keep down the weeds in and around your garden are mulching and cultivation. If proper attention is given to this problem early much time and effort can be saved. Small weeds are easier to control than large ones. When weeds get started they can cause many headaches and backaches, and retard plant growth.

Mulching is covering the soil around your vegetables with a protective material. Besides controlling weeds, the mulch will conserve moisture, regulate the soil temperature and keep the vegetables cleaner. With mulch very little cultivation is needed. Mulch materials include leaves, straw, sawdust, wood chips, cardboard, newspaper, shredded paper, old carpet, and paper and plastic sheeting. On established plantings, materials are spread around the plants. With paper or plastic sheeting the material is rolled out on the prepared seedbed and anchored on the edges with soil. Seeds and transplants are planted through holes at the desired spacings. Water can be applied from the side through furrow irrigation or by a trickle/drip tube or soaker hose under the mulch.

Cultivate with a sharp hoe or cultivator just as the weeds begin to sprout. Scrape and loosen the total soil surface around the plants without going too deep, which would cut or damage shallow roots of the vegetable plants. Cultivation will also help aerate the soil and can be used to mix a side-dressing of nitrogen fertilizer into the soil.

Chemical herbicides for weed control are not generally recommended for use in home gardens.

Be prepared for pests and problems

Problems of the garden can be minimized by being prepared for them. Learn about the insects and diseases that commonly occur in the area and learn control methods. Whenever possible select disease resistant varieties. Soil problems can be reduced if the steps mentioned earlier are followed; however, crop injury from salt can appear if proper management has not been followed. High temperature and shallow watering often cause problems especially when plantings are made too late in the spring or too early in the fall. Also, as temperatures increase more pest problems will occur; be prepared for them. Learn as much as possible from books, bulletins and professionals. Experience is the best teacher on how to handle these problems. Recording treatments in a gardening notebook will be helpful in the future when they occur again.

Harvest at peak quality

The job is not done until top quality vegetables are harvested from the garden. When the "fruits" of your labor are tasted, then it will be worth all the effort.

Most vegetables are at peak quality for only a short period of time and should be harvested. Learn to tell the proper time to harvest each crop. Immature vegetables will not improve after harvest and overmature vegetables will be tough and lack the desired taste and texture. To maintain quality after harvest, handle vegetables carefully. Cool and store vegetables like asparagus, broccoli, leafy crops, peas and sweet corn below 40° F.; tomatoes, peppers, cucumbers and eggplant around 55° F. Remove "field heat" as soon as possible by placing them in the shade or a refridgerator, unless they are eaten immediately.

Garden vegetables offer you a variety of experiences and flavors throughout the year. Enjoy them both.

Elevations of locations in Arizona (feet above sea level)							
10 to 1000'	Feet	1000 to 2000'	Feet	2000 to 3000'	Feet	3000 to 4500'	Feet
Buckeye	888	Ajo	1763	Mammoth	2348	Benson	3585
Gila Bend	737	Casa Gd.	1390	Roosevelt	2200	Bowie	3765
Mohawk	538	Chandler	1213	Safford	2920	Camp Verde	3160
Parker	425	Florence	1500	San Carlos	2630	Chino Valley	4250
Quartzsite	875	Mesa	1225	Sells	2375	Clarkdale	3550
Wellton	260	Phoenix	1108	Superior	2820	Clifton	3465
Yuma	141	Red Rock	1864	Tucson	2423	Cottonwood	3550
Yuma Mesa	181	Salome	1700	Wickenburg	2093	Douglas	3973
3000 to 4500'	Feet	4500 to 6000'	Feet	4500 to 6000'	Feet	6000' and above	Feet
Duncan	3535	Ashfork	5140	St. Johns	5560	Alpine	8000
Globe	3540	Bisbee	5350	Sierra Vista	4620	Flagstaff	6993
Kingman	3333	Chinle	5538	Snowflake	5644	Fort Valley	7347
Nogales	3865	Colorado City	4980	Sonoita	4865	Grand Canyon	6890
Page	4380	Fredonia	5000	Tombstone	4540	Heber	6439
Patagonia	4044	Holbrook	5075	Whiteriver	5280	Pinedale	6500
San Simon	3613	Jerome	5245	Winslow	4850	Show Low	6331
Sedona	4240	Payson	4930	Young	5577	Springerville	6964
Willcox	4182	Prescott	5354	Kayenta	5798	Window Rock	6750

When to plant vegetables in the Arizona garden

Vegetables differ in their climatic requirements making it necessary to know when to plant them in order to have a successful garden.

Some vegetables will withstand cool and even freezing weather. Others need warmer conditions to germinate and to produce. Generally vegetables are placed in two categories—cool-season crops and warm-season crops.

Cool-season vegetables include beet, broccoli, cabbage, carrot, lettuce, onion, pea, potato, radish, spinach and turnip. These are hardy or frost tolerant plants and germinate in cold soil. They can be planted in the fall, winter or early spring depending on location. For best quality these crops need to mature during cooler periods rather than in the heat of the summer.

Warm-season vegetables include beans, cucumber, eggplant, melons, pepper, pumpkin, squash, sweet corn, sweet potato and tomato. These do not tolerate frost but need warm temperatures to set and properly mature fruit. However, high temperatures reduce quality and cause flower abortion. Examples: sunburned fruit, poorly colored tomatoes, poor ear fill of sweet corn, and flowers of tomato, pepper and beans drop. Elevation is indicative of climate. In Arizona gardening occurs from almost sea level to over 7,000 feet. Two problem periods exist—the hot summer at lower elevations and cold winter at higher elevations. Since these conditions should be avoided for many vegetables, considerations should be made when planning the garden planting schedule.

At lower elevations up to 3,000 feet, two main planting periods are generally followed—early spring period for warm-season vegetables and late summer to winter period for cool-season crops. In the higher elevations 3,000 to 6,000 feet, there is one main cropping period which is planted during the spring and early summer. Although, at these elevations in Central and Southern Arizona, an early fall planting of cool season vegetables is usually productive.

The lists below give suggested planting dates for different elevations. These guides are based on experience, observation, frost dates, hardiness and other characteristics of vegetable species. Elevations for certain locations in Arizona are listed above. Find the elevation closest to your location and use these dates along with local experience to develop a vegetable planting program. County Cooperative Extension Offices can offer advice as well as local nurseries and garden centers. These diagrams show some commonly-used systems for growing garden vegetables.



Figure A: The soil-bed technique allows for furrow irrigation water to move from furrow ditches into the bed, pushing salts to the center. To avoid salt problems plant near the bed edge. When using furrow irrigation a slight slope is needed so water will run down the furrow.



Figure B: Make raised beds using railroad ties, landscaping wood, lumber, blocks or rocks. The bed is filled with at least one foot of soil, organic matter, sand, perilite and other materials that promote good plant growth. Raised beds should be used when an area does not have good soil.

In windy areas, sunken beds might be considered to protect young plants and collect water.



Figure C: Water can be applied by drip or soaker hose, furrow, or sprinkler, irrigation. There are many types of systems available that apply water efficiently. Whatever method is used, adequate watering moves salts down and away from the plant roots. Select a system that meets the need and can be managed properly.

VEGETABLE PLANTING DATES

Vegetable Species	10 to 1000 feet	1000 to 2000 feet	2000 to 3000 feet		
Asparagus	Oct. 1-Feb. 1	Oct. 1-Mar. 1	Oct. 1-Mar. 1		
Bean,bush	Feb. 1-Mar. 1	Feb. 15-Mar. 15	Mar. 1-Apr. 1		
	Aug. 1-Sept. 1	July 25-Aug. 15	July 15-Aug. 15		
Bean, pole	Aug. 1-Sept. 1	July 15-Aug. 15	July 15-Aug. 10		
Bean, lima	Feb. 1-Mar. 1	Feb. 15-Mar. 15	Mar. 1-Apr. 1		
Bean, edible soy	Mar. 1-May 1	Mar. 15-June 1	Apr. 1-June 1		
Beet	Sept. 15-Mar. 1	Sept. 1-Mar. 15	Aug. 25-Apr. 1		
Broccoli	Sept. 1-Jan. 1	Sept. 1-Dec. 1	July 25-Oct. 1		
Brussels Sprouts	Sept. 1-Jan. 1	Sept. 1-Dec. 1	Aug. 15-Oct. 1		
Cabbage (seed)	Sept. 1-Nov. 20	Aug. 15-Dec. 1	Aug. 1-Dec. 1		
Cabbage (plants)	Oct. 1-Dec. 1	Sept. 15-Jan. 1	Sept. 1-Feb. 1		
Cantaloupe	Dec. 1-Apr. 10	Feb. 15-Apr. 1	Mar. 15-June 1		
Carrot	July 15-Aug. 15	Sept. 1-Mar. 1	Aug. 25- Mar.15		
	Sept. 1-Jan. 1				
Cauliflower	Same as cabbage	Same as cabbage	Same as cabbage		
Celery	October 15	Aug. 15-Oct. 15	Aug. 1-Oct. 15		
Chard	Sept. 1-Jan. 1	Sept. 1-Mar. 1	Aug. 15-Apr. 1		
Chinese Cabbage	Sept. 15-Dec. 1	Sept 1-Jan. 1	Aug.15-Jan.15		
Collard	Sept. 15-Dec. 1	Sept. 1-Jan. 1	Sept.1 -Jan.15		
Corn, sweet	Feb. 15-Mar. 1	Feb. 15-Mar. 15	Mar. 15-Apr. 1		
	July 30-Aug. 30	July 20-Aug. 20	July 15-Aug. 15		
Corn, Mexican June		June 20-July 20	July 1-July 5		
Cucumber	Dec. 1-Apr. 1	Mar. 1-Apr. 1	Mar. 20-May 15		
		Aug. 15-Sept. 15	Aug. 1-Sept. 1		
Eggplant	Jan. 15-Apr. 1	Feb. 1-Apr. 1	Apr. 1-May 15		
Endive	Sept. 1-Dec. 1	Sept. 1-Jan. 1	Sept. 1-Feb. 1		
Garlic	Sept. 1-Dec. 1	Sept. 1-Dec. 1	Sept. 1-Jan. 1		
Horseradish	Not adapted	Not adapted	Nov. 1-Feb. 1		
Kale	Sept. 1-Dec. 1	Sept. 1-Dec. 1	Aug. 15-Feb. 15		
Kohlrabi	Sept. 1-Dec. 1	Sept. 1-Dec. 1	Sept. 1-Feb. 1		
Leek	Sept. 15-Dec. 15	Sept. 1-Jan. 1	Sept. 1-Jan. 15		
Lettuce, head	Sept. 20-Nov. 20	Sept. 1-Jan. 1	Sept. 1-Feb. 15		
Lettuce, leaf	Sept. 20-Jan. 1	Sept. 1-Mar. 1	Aug. 20-Apr. 1		
Muskmelon	Dec. 1-Apr. 10	Feb. 15-Apr. 1	Apr. 1-July 15		
		July 1-Aug. 1			
Mustard	Sept. 15-Dec. 15	Sept. 1-Jan. 1	Sept. 1-Feb. 1		
Okra	Mar. 1-Apr. 15	Mar. 1-June 1	AprJune 15		
Onion, green bunch	Sept. 15-Jan. 15	Sept. 1-Feb. 1	Aug. 15-Feb. 1		
Onion, dry (seeds)	Nov.1-Dec.15	Oct. 15-Jan. 1	Oct. 15-Jan. 1		
Onion, dry (sets)	Nov. 15-Jan. 15	Nov. 1-Feb. 1	Nov. 1-Feb. 15		

Vegetable Species	10 to 1000 feet	1000 to 2000 feet	2000 to 3000 feet	
Parsley	Oct. 1-Jan. 15	Sept. 1-Jan. 1	Sept. 1-Jan. 15	
Parsnip	Not adapted	Sept. 1-Jan. 1	Sept. 1-Jan. 15	
Pea, fall	Sept. 10-Sept. 20	Aug. 15-Sept. 15	Aug. 15-Sept. 15	
Pea, spring	Jan. 20-Feb. 15	Oct. 15-Dec. 15	Feb. 1-Mar. 15	
Pepper (seed)	NovJan.	Feb. 1-Mar. 1	Feb. 15-Mar. 15	
Pepper (plants)	Feb. 1-Mar. 15	Mar. 1-Apr. 1	Apr. 1-June 1	
Potato, Irish	Sept. 1-Feb. 15	Feb. 1-Mar. 15	Feb. 15-May 1	
Potato, sweet	Mar. 1-June 20	Mar. 1-June 1	May 1-June 15	
Pumpkin	July 15-Aug. 15	July 1-Aug. 1	Apr. 1-July 15	
Radish	Sept. 1-Apr. 1	Sept 1-Apr. 15	Aug. 5-May 1	
Rhubarb	Not adapted	Not adapted	Oct. 1-Mar. 1	
Rutabaga	Sept. 15-Jan. 15	Sept. 1-Feb. 1	Aug. 20-Mar. 1	
Salsify	Not adapted	Not adapted	Oct. 1-Dec. 1	
Spinach	Sept. 15-Feb. 1	Sept. 1-Feb. 1	Aug. 20-Mar. 1	
Squash, summer	Dec. 15-Apr. 10	Feb. 1-May 1	Mar. 15-July 15	
Squash, winter	July 15-Aug. 15	July 1-31	July 1- July 31	
Tomato (seed)	NovJan.	Jan. 1-Mar.1	Jan. 10-Feb. 15	
Tomato (plants)	JanMar. 15	Feb. 15-Mar. 15	Mar. 15- Apr. 15	
Turnip	Sept. 15-Feb. 1	Sept. 1-Feb. 1	Aug. 15- Mar. 1	
Watermelon	Dec. 15-Apr. 1	Feb. 15-Apr. 1	Mar. 15- June 1	
Vegetable Species	3000 to 4500 feet	4500 to 6000 feet	Above 6000 feet	
Asparagus	Feb. 15-Apr. 1	April 1-30	Apr. 15-May 15	
Bean, bush	Apr. 25-July 15	May 15-July 1	May 25-June 15	
Bean, pole	Apr. 25-July 15	May 15-July 1	May 25-June 15	
Bean, lima	Apr. 25-July 15	May 15-July 1	May 25-June 15	
Bean, edible soy	May 15-July 1	May 25-July 1	Not adapted	
Beet	Mar. 1-May 15	May 1-July 15	May15-June15	
Broccoli	Feb. 15-Aug. 1	Apr. 1-July 15	Not adapted	
	Sept. 1-Oct. 15			
Brussels Sprouts	July 1-Aug. 1	June 1-July 1	May 15-June 15	
Cabbage (seed)	Feb. 15-Apr. 15	March 15-30	April 1-15	
Cabbage (plants)	Mar. 15-May 1			
	Aug. 20-Oct. 1	May 1- June 1	May 15-June 15	
Cantaloupe	May 1-June 20	May 15-June 15	May 25-June 10	
Carrot	Mar. 1-May 10	May 1-July 15	May 15-July 1	
	July 15-Sept. 15			
Cauliflower	Same as cabbage	Same as cabbage	Same as cabbage	
Celery (plants)	May 15-June 20	June 1-July 15	Not adapted	
Chard	July 15-Sept. 15	July 1-Aug. 1		
	Feb. 15-Apr. 30	Mar. 1-Apr. 10	Apr. 1-June 10	
Chinese Cabbage	July 1-Sept.15	June 1-July 15	May 15-June 15	
Collard	June 15-Aug. 1	June 1-July 15	May 15-July 1	

Vegetable Species	3000 to 4500 feet	4500 to 6000 feet	Above 6000 feet	
Corn, sweet	May 10-July 15	May 25-July 1	June 1-10	
Corn, Mexican June	May 10-July 15	May 25-June 15	Not adapted	
Cucumber	May 10-June 15	May 15-June 15	June 1-25	
Eggplant (plants)	May 1-June 15	May 15-June 15	June 1-20	
Endive	Feb. 1-Apr. 1	Apr. 15-June 15	May 15-June 15	
Garlic	Feb. 15-Apr. 10	April 1-30 (cloves)	Not adapted	
	Sept. 15-Nov. 15			
Horseradish	FebApr.	Feb. 15-Mar. 15	April-May	
Kale	Feb. 1-Mar. 20	Feb. 15-Apr 10	April-May	
	Aug. 1-Sept. 15			
Kohlrabi	Feb. 15-Apr. 1	Apr.15-May15	May 15-June 1	
Leek	Feb. 15-Apr. 10	April 1-30	Not adapted	
Lettuce, head	Feb. 15-Mar. 15	July 1-Aug. 1	June 1-30	
	July 15-Aug. 15			
Lettuce, leaf	Mar. 1-Apr. 15	Mar. 15-Apr.15	May 1- July 1	
	July 15-Sept. 15	Aug. 1-Sept.15		
Muskmelon	May 10-June 15	May 15-June 15	Not adapted	
Mustard	Feb. 15-July 15	Apr. 1-July 1	April-May	
Okra	May 10-July 1	May 15-June 15	June 1-10	
Onion, green bunch	Feb. 15-May 1	Apr. 15-May 1	May 1-31	
Onion, dry (seeds)	Jan. 15-Mar. 15	Feb. 15-Apr. 15	April 1-30	
	Sept. 15-Nov. 15	Oct. 15-Jan. 1	Oct. 15-Jan. 1	
Onions, dry (sets)	Sept. 15-Nov. 15	Nov. 1-Feb. 1	Nov. 1-Feb. 15	
	Feb. 15-Apr. 15	Apr. 1-15	Apr. 15-June 1	
Parsley	May 1-June 15	Apr. 1-15	May 1-31	
Parsnip	Mar. 1-May 1	Apr. 1-May 20	April-May	
Pea, spring	Feb. 1-Mar. 15	Feb. 15-Aug. 15	May 1-June 1	
Pea, fall	Aug. 25-Oct. 15	Aug. 1-Sept. 1	Not adapted	
Pepper (seed)	Feb. 15-Mar. 30	Mar. 1-Apr. 1	Apr. 1-15	
Pepper (plants)	May 10-June 1	May 10-Aug. 25	May 15-June 1	
Potato, Irish	Mar. 20-Apr. 20	May 10-June 1	May 15-June 1	
	July 25-Aug. 15			
Potato, sweet	May 10-25	May 15-20	Not adapted	
Pumpkin	May 15-July 1	May 20-June 15	May 25-June 10	
Radish	Mar. 1-May 15	Apr. 1-June 15	May 15-June15	
	July 15-Sept. 15		,	
Rhubarb	Mar. 1-Apr. 20	Mar. 1-Apr. 1	April 1-30	
Rutabaga	Mar. 1-Apr. 1	Apr. 1-May 15	May 1-June 1	
Salsify	Mar. 15-June 1	Apr. 1-May 15	May 1-June 1	
Spinach	Feb. 15-Apr. 15	Apr. 1-May 15	May 1-June 1	
1	Sept. 15-Oct. 15	P		
Squash. summer	May 10-July 15	May 1-July 1	May 15-June 15	
Squash, winter	May 10-July 1	May 15-July 1	May 15-June 10	

Vegetable Species	3000 to 4500 feet	4500 to 6000 feet	Above 6000 feet
Tomato (seed)	Mar. 1-Apr. 1	Mar. 1-Apr. 1	Apr. 1-10
Tomato (plants)	May 1-June 15	May 10-June 1	May 25-June 10
Turnip	Mar. 1-Apr. 15	Apr. 1-May15	May 15-June 1
	Aug. 15-Sept.15		
Watermelon	May 10-July 15	May 1-June 1	Not adapted



THE UNIVERSITY OF ARIZONA COLLEGE OF AGRICULTURE AND LIFE SCIENCES TUCSON, ARIZONA 85721

Norman F. OEBKER Vegetable Specialist, Emeritus

ROBERT E. CALL Horticulture Agent

CONTACT: ROBERT E. CALL recall@ag.arizona.edu

This information has been reviewed by university faculty. cals.arizona.edu/pubs/garden/az1435.pdf

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, James A. Christenson, Director, Cooperative Extension, College of Agriculture & Life Sciences, The University of Arizona.

The University of Arizona is an equal opportunity, affirmative action institution. The University does not discriminate on the basis of race, color, religion, sex, national origin, age, disability, veteran status, or sexual orientation in its programs and activities.